

**General characteristics of multi-partite
quantum systems**
**(Lecture of the Quantum Information class of
the Master in Quantum Science and
Technology)**

Géza Tóth

Theoretical Physics, University of the Basque Country (UPV/EHU), Bilbao, Spain
Donostia International Physics Center (DIPC), San Sebastián, Spain
IKERBASQUE, Basque Foundation for Science, Bilbao, Spain
Wigner Research Centre for Physics, Budapest, Hungary

UPV/EHU, Leioa
19, 21 and 16 January, 2021

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

A single classical bit

- A classical bit can be either 0 or 1. Can we still use it to describe a real number between 0 and 1?
- For that, we need an ensemble of several classical bits

$$\{b_k\}_{k=1}^M, \quad (1)$$

where $b_k = 0$ or 1

- We can interpret the average value and the variance. That is,

$$\langle b \rangle = \frac{1}{M} \sum_k b_k, \quad (2)$$

and

$$(\Delta b)^2 = \frac{1}{M} \sum_k (b_k - \langle b \rangle)^2. \quad (3)$$

A single classical bit II

- This can also be given with probabilities:
- Let P_0 and P_1 be the probabilities of having a 0 or a 1.
- The expectation value and the variance are the function of P_0 and P_1 . Since $P_0 + P_1 = 1$, we have a **single real degree of freedom** that describes the statistical properties of an ensemble of bits.
- Hence,

$$\langle b \rangle = P_1 \quad (4)$$

and

$$(\Delta b)^2 = P_0(0 - P_1)^2 + P_1(1 - P_1)^2. \quad (5)$$

Stochastic computing

- Stochastic computing uses random bits to calculate (John von Neumann, 1953).
- A random bit represents a real number between 0 and 1. Two random bits can easily be multiplied.

$$\langle b_1 b_2 \rangle = \langle b_1 \rangle \langle b_2 \rangle. \quad (6)$$

- We need many samples to get the average with small error.

Several classical bits

- N classical bits can be in one of the 2^N binary states. For example, for $N = 2$, these are 00, 01, 10 and 11.
- The ensemble of the N -bit units can be described by the 2^N probabilities.
- For $N = 2$, these are

$$P_{00}, P_{01}, P_{10}, P_{11}. \quad (7)$$

- Since, again, the sum of all the probabilities is 1, **we need** $2^N - 1$ **real degrees of freedom** to describe the statistical properties of such an ensemble.

Several classical bits II

- Let us consider some function of N bits $f(k)$, where k is now an N bit number.
- Then, the expectation value of f is

$$\langle f \rangle = \sum_{k=0}^{2^N-1} p_k f(k) = \vec{p} \vec{f}, \quad (8)$$

where k is an N -bit number, i.e., an integer between 0 and $2^N - 1$. We put the f_k 's into a vector \vec{f} . We also put the p_k probabilities into \vec{p} .

Several classical bits III

- We can also write

$$\langle f^2 \rangle = \sum_k p_k f_k^2 \quad (9)$$

Hence,

$$(\Delta f)^2 = \sum_k p_k f_k^2 - \left(\sum_k p_k f_k \right)^2. \quad (10)$$

These were relevant, since in the quantum case, we will have similar expressions.

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Quantum bit - pure states

- A quantum bit (=two-state system, spin- $\frac{1}{2}$ particle) can be in a pure state

$$|q\rangle = \alpha|0\rangle + \beta|1\rangle, \quad (11)$$

where α and β are complex numbers, and the normalisation condition $\alpha^2 + \beta^2 = 1$.

- Note that the overall phase does not matter, thus a pure quantum bit is described by two degrees of freedom. (The two complex coefficients have **4 real degrees of freedom**.)
- However, due to the normalisation condition and the arbitrariness of the overall phase we are left with **two degrees of freedom**.)

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- **C. Multi-qubit systems - pure states**
- D. Measurement
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Multi-qubit systems - pure states

- What about a two-qubit system? What kind of states it can be in?
One could think on qubit 1 in state

$$|q_1\rangle = \alpha_1|0\rangle + \beta_1|1\rangle, \quad (12)$$

and qubit 2 in state

$$|q_2\rangle = \alpha_2|0\rangle + \beta_2|1\rangle. \quad (13)$$

- However, we all know that the general state of the two-qubit system can be given as

$$|q_{12}\rangle = \alpha_{00}|00\rangle + \alpha_{01}|01\rangle + \alpha_{10}|10\rangle + \alpha_{11}|11\rangle. \quad (14)$$

Multi-qubit systems - pure states II

- In general, for N qubits we need N complex numbers. Again the state has to be normalized and the overall phase does not matter, thus this means $2 \times 2^N - 2$ real degrees of freedom.
- We can place the coefficients in a vector, called state vector and write

$$|q_{12}\rangle = \begin{pmatrix} \alpha_{00} \\ \alpha_{01} \\ \alpha_{10} \\ \alpha_{11} \end{pmatrix}. \quad (15)$$

- The properties of the state vector are: it is normalized

$$\langle \Psi | \Psi \rangle = 1. \quad (16)$$

Multi-qubit systems - pure states III

- An overall phase does not matter:

$$e^{-i\theta}|\Psi\rangle \quad (17)$$

describes the same state for any θ .

- The expectation value of an operator for a pure state can be obtained as

$$\langle A \rangle = \langle \Psi | A | \Psi \rangle = \text{Tr}(A |\Psi\rangle \langle \Psi|). \quad (18)$$

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- **D. Measurement**
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Measurement

- The von Neumann measurement in the z basis results is either 0 or 1. If the state was Eq. (11), then we get a statistical mixture of 0 and 1, with the probabilities

$$P_0 = |\alpha|^2, \quad (19)$$

and

$$P_1 = |\beta|^2. \quad (20)$$

That is, from an ensemble of quantum bits we get an ensemble of classical bits.

- If we measure in the x basis, we get another classical ensemble.
- For a multi-qubit system, if we measure in the some basis (e.g., x , y or z), we get an ensemble of N -bit systems. However, for each choice of basis we get a different classical ensemble.

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- **E. Mixed states and the density matrix**
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Mixed states and the density matrix

- So far we were talking about pure states.
- In reality, in an experiment we do not have a situation where a machine always produces the $|\Psi_1\rangle$ state.
- Sometimes it makes mistakes, and produces the $|\Psi_k\rangle$ states for $k = 2, 3, \dots$ How to describe such a situation?

$ \Psi_1\rangle$	p_1
$ \Psi_2\rangle$	p_2
$ \Psi_3\rangle$	p_3
...	...

Mixed states and the density matrix

- What is the expectation value of an operator in such a system?
We can write it as

$$\langle A \rangle = \sum_k p_k \langle \psi_k | A | \psi_k \rangle = \text{Tr} \left(A \sum_k p_k | \psi_k \rangle \langle \psi_k | \right). \quad (21)$$

- This can be rewritten as

$$\langle A \rangle = \text{Tr}(\rho A), \quad (22)$$

where

$$\rho = \sum_k p_k | \psi_k \rangle \langle \psi_k | \quad (23)$$

is the density matrix (Neumann, Landau).

- Note that if ρ is diagonal, we obtain

$$\langle A \rangle = \text{Tr}(\rho A) = \sum_k \rho_{kk} A_{kk}. \quad (24)$$

That is, A is written in the eigenbasis of ρ . This is the scalar product of two vectors as in $\langle f \rangle = \vec{p} \vec{f}$ [See Eq. (8)].

Mixed states and the density matrix II

- The density matrix describes the state completely. Now we see, why the overall phase does not matter:

$$e^{-i\theta}|\Psi_k\rangle\langle\Psi_k|e^{+i\theta} = |\Psi_k\rangle\langle\Psi_k|. \quad (25)$$

- The properties of the density matrix are

$$\begin{aligned} \rho &= \rho^\dagger, \\ \rho &\geq 0, \\ \text{Tr}(\rho) &= 1. \end{aligned} \quad (26)$$

- **A $2^N \times 2^N$ density matrix has $4^N - 1$ real parameters.**
- For $N = 1$, this means 3 real parameters, corresponding to the three coordinates of the Bloch vector. For $N = 2$, this means 8 real parameters.

Mixed states and the density matrix III

- We can also say that

$$\text{Tr}(\rho^2) \leq 1. \quad (27)$$

It is one only for pure (rank-1) states.

- The density matrix can be decomposed into the sum of pure states in many ways. The decomposition

$$\rho = \sum_k p_k |\psi_k\rangle\langle\psi_k| \quad (28)$$

is not unique, i.e., it is not necessarily an eigendecomposition. This has a large importance for entanglement theory.

Mixed states and the density matrix IV

Summary:

	N bits	N qubits
Number of DOF	$2^N - 1$	$4^N - 1$
Description	\vec{p}	ρ
Expectation value	$\vec{f}\vec{p}$	$\text{Tr}(A\rho)$
Normalization	$\sum_k p_k = 1$	$\text{Tr}(\rho) = 1$

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- E. Mixed states and the density matrix
- **F. Geometry of quantum states**
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Bloch vector

- For a single qubit, **the density matrix has three real parameters**. It can be written as

$$\varrho = \frac{1}{2} \left(\mathbb{1} + \sum_{l=x,y,z} v_l \sigma_l \right), \quad (29)$$

where σ_l are the Pauli spin matrices.

- Using $\text{Tr}(\sigma_k \sigma_l) = 2\delta_{kl}$, we can write

$$\text{Tr}(\varrho^2) = \frac{1}{2} + \frac{1}{2} \sum_{l=x,y,z} v_l^2. \quad (30)$$

That is, the Bloch vector has a maximal length for pure states.

Bloch vector II


- From $\text{Tr}(\rho^2) \leq 1$, the condition for being physical is Eq. (26), which is equivalent to

$$\sum_{l=x,y,z} |v_l|^2 \leq 1. \quad (31)$$

The three-element vector is called the Bloch vector.

Bloch vector III

- Let us identify the points in (v_x, v_y, v_z) corresponding to physical states. They are in a ball.
- The pure states are on the surface.
- Mixed states are inside the Ball. This is because $\text{Tr}(\rho^2)$ is directly related to the length of the Bloch vector.
- The $|0\rangle$ and $|1\rangle$ correspond to the North and South Pole.
- $|0\rangle + \exp(-i\phi)|1\rangle$ correspond to points on the equator. See Fig. ??



Set of physical quantum states for a single qubit. The axes correspond to v_l for $l = x, y, z$. Pure states correspond to points on the surface, mixed states correspond to internal points.

A single qudit (qunit): d -dimensional systems

- For higher dimensional systems the picture is much more complicated. Let us consider qudits with dimension d .
- Similarly to the case before, a $d \times d$ **Hermitian matrix with a unit trace has $d^2 - 1$ degrees of freedom**. Hence, we can write a density matrix as a linear combination of $d^2 - 1$ SU(d) generators as

$$\rho = \frac{1}{d} \mathbb{1} + \frac{1}{2} \sum_{l=1}^{d^2-1} v_l g_l. \quad (32)$$

Here,

$$\text{Tr}(g_k g_l) = 2\delta_{kl}. \quad (33)$$

(Like for the Pauli matrices. Thus, we have something like the generalized Pauli matrices. $d = 3$: Gell-Mann matrices.)

A single qudit (qunit): d -dimensional systems II

- One can check that

$$\text{Tr}(\rho g_I) = v_I. \quad (34)$$

- With these, we can write

$$\text{Tr}(\rho^2) = \frac{1}{d} + \frac{1}{2} \sum_{I=1}^{d^2-1} v_I^2. \quad (35)$$

- For physical states the purity is not larger than one. Hence, the Bloch vector still has a length limit


$$\sum_{I=1}^{d^2-1} |v_I|^2 \leq L_{\max}^2 = 2 \frac{d-1}{d}. \quad (36)$$

- However, not all values satisfying this length constraint correspond to some physical quantum state. Again, for pure states the Bloch vector has maximal length. Thus pure states are on the sphere. Note also that for finite d the purity cannot be zero.

A single qudit (qunit): d -dimensional systems III

- Let us again look at the points $(v_1, v_2, \dots, v_{d^2-1})$ corresponding to physical states.
- First note that the set is convex. This is because mixing two physical states we always get a physical state.


A single qudit (qunit): d -dimensional systems IV



Two convex and a concave object.

A single qudit (qunit): d -dimensional systems V

- On the next figure we will show the set of quantum states. The coordinate axis could be the v_i , for example. Inside the set there are the density matrices with full rank. On the boundary there are the states with less than full rank, such as for example rank-1 states, which are pure states.



Set of physical quantum states. Note that the set is convex.
A,B,D: rank-1 states. C: rank-2 state. E: full rank states.

A single qudit (qunit): d -dimensional systems VI

- **Observation.** The following inequality is true

$$\lambda_{\min}(\mathbf{A} + \mathbf{B}) \geq \lambda_{\min}(\mathbf{A}) + \lambda_{\min}(\mathbf{B}). \quad (37)$$

Proof. Let us consider that for a Hermitian matrix X we have

$$\lambda_{\min}(X) = \min_{\psi} \langle \psi | X | \psi \rangle. \quad (38)$$

Then, for A and B Hermitian matrices we have

$$\lambda_{\min}(\mathbf{A} + \mathbf{B}) = \min_{\psi} \langle \psi | \mathbf{A} + \mathbf{B} | \psi \rangle \geq \min_{\psi} \langle \psi | \mathbf{A} | \psi \rangle + \min_{\psi} \langle \psi | \mathbf{B} | \psi \rangle = \lambda_{\min}(\mathbf{A}) + \lambda_{\min}(\mathbf{B}) \quad (39)$$

□

We can prove similarly that

$$\lambda_{\max}(\mathbf{A} + \mathbf{B}) \leq \lambda_{\max}(\mathbf{A}) + \lambda_{\max}(\mathbf{B}). \quad (40)$$


Full rank states

- Using this, we can say the following.
- **Observation.** Full-rank states are inside the set.
Proof. If the state is full rank, it means that for some small ϵ

$$\varrho' = \varrho + \epsilon H \quad (41)$$

is also physical, where H is a trace 0 Hermitian matrix. Why is that? See also the next figure.

Full rank states II



We take an internal state ρ and consider the states ρ' in its neighborhood.

Full rank states III

- It is physical since
 - 1 Trace is 1.
 - 2 Hermitian.
 - 3 Eigenvalues are nonzero for small epsilon. This is because

$$\lambda_{\max}(\rho) + \lambda_{\max}(\epsilon H) \geq \lambda_k(\rho') \geq \lambda_{\min}(\rho) + \lambda_{\min}(\epsilon H). \quad (42)$$

Here we have

$$\lambda_{\min}(\epsilon H) = \begin{cases} +\epsilon \lambda_{\min}(H), & \text{if } \epsilon \geq 0, \\ -\epsilon \lambda_{\max}(H), & \text{if } \epsilon < 0. \end{cases} \quad (43)$$

Similar statement holds for $\lambda_{\max}(\epsilon H)$. \square

Non-full-rank states

- **Observation.** Non-full-rank states are on the surface of the set.
- *Proof.* If the state is not full rank, then it has zero eigenvalues. Thus, there is an H such that ϱ' is aphysical for any $\epsilon > 0$ or any $\epsilon < 0$.
- To be more explicit, let us write

$$\varrho = UDU^\dagger, \quad (44)$$

such that D contains the eigenvalues. Here,

$$D = \text{diag}(\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_d), \quad (45)$$

and the eigenvectors are

$$U = [|\Psi_1\rangle, |\Psi_2\rangle, |\Psi_3\rangle, \dots, |\Psi_d\rangle]. \quad (46)$$

Non-full-rank states II

- Assume that $\lambda_d = 0$. Then,

$$\varrho' = \varrho + \epsilon(|\Psi_d\rangle\langle\Psi_d| - \mathbb{1}/d) \quad (47)$$

has a negative eigenvalue for any $\epsilon < 0$. The Identity is needed to make the expression zero-trace.

- This is because the eigenvalues of this matrix are

$$D' = \text{diag}(\lambda_1 - \epsilon, \lambda_2 - \epsilon, \lambda_3 - \epsilon, \dots, \lambda_d + \epsilon(1 - 1/d)). \quad (48)$$

□

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Two or more qubits: reduced states

- How can one see the state of a qubit, if it is the part of an entangled state?
- A reduced state of a bipartite system can be obtained after tracing out one of the subsystems. Let us consider a two-qubit system and write the density matrix in the basis $|00\rangle, |01\rangle, |10\rangle, |11\rangle$. Then, denote the elements of the density matrix by

$$\rho_{ij,kl}, \quad (49)$$

where $i, j, k, l = 0, 1$. In other words, it looks like

$$\rho = \begin{array}{l} |00\rangle \\ |01\rangle \\ |10\rangle \\ |11\rangle \end{array} \begin{pmatrix} \rho_{00,00} & \rho_{00,01} & \rho_{00,10} & \rho_{00,11} \\ \rho_{01,00} & \rho_{01,01} & \rho_{01,10} & \rho_{01,11} \\ \rho_{10,00} & \rho_{10,01} & \rho_{10,10} & \rho_{10,11} \\ \rho_{11,00} & \rho_{11,01} & \rho_{11,10} & \rho_{11,11} \end{pmatrix}. \quad (50)$$

Thus, the size of the density matrix is 4×4 .

Two or more qubits: reduced states II

- To become familiar with bras and kets, one can even use the completeness relation

$$\text{Identity} = \sum_{ij} |ij\rangle\langle ij|. \quad (51)$$

Then, one obtains

$$\text{Identity} \times \rho \times \text{Identity} = \sum_{ijkl} |ij\rangle(\langle ij|\rho|kl\rangle)\langle kl|, \quad (52)$$

where the expression in the bracket is just the matrix element of the density matrix

$$\rho_{ij,kl} = \langle ij|\rho|kl\rangle. \quad (53)$$

Hence, the density matrix can be written as

$$\rho = \sum_{ijkl} \rho_{ij,kl} |ij\rangle\langle kl|. \quad (54)$$

Two or more qubits: reduced states III

- Then, tracing out the second subsystem gives the reduced state

$$\text{Tr}_2(\rho) = \rho_{\text{red}}, \quad (55)$$

which is given as

$$\rho_{\text{red},ik} = \sum_m \rho_{im,km}. \quad (56)$$

This is a 2x2 density matrix of a qubit. With this

$$\langle \mathbf{A} \otimes \mathbb{1} \rangle_\rho = \langle \mathbf{A} \rangle_{\rho_{\text{red}}}. \quad (57)$$

- Graphical representation: in the blockdiagonal representation, we sum the matrices in the diagonal.
- Tracing out for pure states:

$$\text{Tr}_2 \left(\sum_k \alpha_k |\psi_k\rangle \langle \phi_k| \right) = \sum_k |\alpha_k|^2 |\psi_k\rangle \langle \psi_k|. \quad (58)$$

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- **H. Purifications**
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Purifications

- The pure state $|\Psi\rangle_{AB}$ state is the purification of the mixed state ρ_A if

$$\text{Tr}_B(|\Psi\rangle_{AB}\langle\Psi|) = \rho_A. \quad (59)$$

Note that $|\Psi\rangle_{AB}$ lives on subsystems A and B , while ρ_A lives on subsystem A only.

- Let us assume that a density matrix is defined as

$$\rho_A = \sum_k p_k |\phi\rangle\langle\phi|_A. \quad (60)$$

- Then, a purification can be a pure state

$$|\Psi\rangle_{AB} = \sum_k \sqrt{p_k} |\phi\rangle_A \otimes |k\rangle_B, \quad (61)$$

where $|k\rangle_B$ denotes an orthonormal basis of the subsystem B .

Purifications II

- If $|\Psi\rangle_{AB}$ is a purification then

$$|\Psi\rangle'_{AB} = \mathbb{1}_A \otimes U_B |\Psi\rangle_{AB}, \quad (62)$$

is also a purification.

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Purity

- Defined as

$$\text{Tr}(\rho^2). \tag{63}$$

- 1 for pure states.
- $1/d$ for the completely mixed state.

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- **J. Entropy**
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Shannon entropy

- There is a source that outputs an integer number between 1 and d .
- The Shannon entropy is given as

$$H = - \sum_{k=1}^d p_k \log p_k. \quad (64)$$

Shannon entropy II

- Properties
 - Classical, not quantum.
 - In information theory, the entropy of a random variable is the average level of "information", "surprise", or "uncertainty" inherent in the variable's possible outcomes (wikipedia). That is, the source can have d possible outputs with some probability. Relation to compression of data.
 - $H = 0$ if $p_1 = 1$, all other $p_k = 0$. $\vec{p} = (1, 0, 0, 0, \dots)$. The output is always the same. No information is provided. Comment: we can show that, using L'Hospital's rule,

$$\lim_{x \rightarrow 0} (x \log x) = \lim_{x \rightarrow 0} \frac{\log x}{1/x} = \lim_{x \rightarrow 0} \frac{1/x}{-1/x^2} = - \lim_{x \rightarrow 0} x = 0. \quad (65)$$

- $H = \log d$ (maximal) if $p_k = \frac{1}{d}$. $\vec{p} = (\frac{1}{d}, \frac{1}{d}, \frac{1}{d}, \frac{1}{d}, \dots)$. All outputs are equally probable, a lot of information is provided.

Von Neumann entropy

- Von Neumann entropy for a quantum state is defined as

$$S(\rho) = -\text{Tr}(\rho \log \rho) \equiv -\langle \log \rho \rangle. \quad (66)$$

- Note: matrix logarithm! It can be written with the eigenvalues of the density matrix as

$$S(\rho) = -\sum_{k=1}^d \lambda_k \log_2 \lambda_k. \quad (67)$$

Von Neumann entropy II

- Properties

- Quantum. "Quantum version" of the Shannon entropy.
- For a pure state we have $\lambda_k = \{1, 0, 0, \dots, 0\}$, and thus it is zero.
- Its maximal is for the completely mixed state for which $\lambda_k = \{\frac{1}{d}, \frac{1}{d}, \frac{1}{d}, \dots, \frac{1}{d}\}$, and its value is $\log_2 d$.
- Concave, i.e., $S(p\rho_1 + (1-p)\rho_2) \geq pS(\rho_1) + (1-p)S(\rho_2)$. Put the figure of a concave function and explain concavity.
- Additive for independent systems, $S(\rho_1 \otimes \rho_2) = S(\rho_1) + S(\rho_2)$.
- Strongly subadditive, $S(\rho_{ABC}) + S(\rho_B) \leq S(\rho_{AB}) + S(\rho_{BC})$. The matrices ρ_B, ρ_{AB} , etc. reduced states.
- Subadditive, $S(\rho_{AC}) \leq S(\rho_A) + S(\rho_C)$.
- Often used in condensed matter physics and field theory. See block entropy depending on the block size.

Quantum relative entropy

- The relative entropy is given as

$$S(\varrho\|\sigma) = -\text{Tr}[\varrho(\log \sigma - \log \varrho)] = -\text{Tr}(\varrho \log \sigma) - S. \quad (68)$$

- Properties

- $S(\varrho\|\sigma) \geq 0$.
- $S(\varrho\|\sigma) = 0$ if and only if $\varrho = \sigma$.
- Not symmetric $S(\varrho\|\sigma) \neq S(\sigma\|\varrho)$.
- Sort of a distance between two quantum states.

Linear entropy

- The linear entropy is defined as

$$S_{\text{lin}}(\rho) = 1 - \text{Tr}(\rho^2) \equiv \langle \mathbb{1} - \rho \rangle. \quad (69)$$

- It is often easier to obtain than the von Neumann entropy.
- Its relation to von Neumann entropy via the Mercator series is

$$-\langle \log \rho \rangle = \langle \mathbb{1} - \rho \rangle + \langle (\mathbb{1} - \rho)^2 \rangle / 2 + \langle (\mathbb{1} - \rho)^3 \rangle / 3 + \dots \quad (70)$$

This is based on expanding

$$\log(\mathbb{1} - (\mathbb{1} - \rho)) \quad (71)$$

Mercator series

$$\log(1 + x) = x - x^2/2 + x^3/3 - + \dots \quad (72)$$

Note that

$$\mathbb{1} - \rho \geq 0. \quad (73)$$

Hence,

$$S \geq S_{\text{lin}}. \quad (74)$$

1 General characteristics of multi-partite quantum systems

- A. Classical bits
- B. Quantum bit - pure states
- C. Multi-qubit systems - pure states
- D. Measurement
- E. Mixed states and the density matrix
- F. Geometry of quantum states
 - A single qubit
 - A single qudit (qunit): d -dimensional systems
- G. Two or more qubits: reduced states
- H. Purifications
- I. Purity
- J. Entropy
 - Shannon entropy
 - Von Neumann entropy
 - Quantum relative entropy
 - Linear entropy
- K. Fidelity

Fidelity

How to measure the distance between quantum states?

- Pure states: overlap square, $\langle \psi | A | \phi \rangle^2$. $\langle \psi | A | \psi \rangle = 0$ if and only if $|\psi\rangle = |\phi\rangle$.
- A pure state and a mixed state:

$$\text{Tr}(|\Psi\rangle\langle\Psi|\varrho) = \langle\Psi|\varrho|\Psi\rangle. \quad (75)$$

- Two mixed states: more difficult

$$F(\varrho, \sigma) = \left(\text{Tr}(\sqrt{\sqrt{\varrho}\sigma\sqrt{\varrho}}) \right)^2. \quad (76)$$

- $0 \leq F(\varrho, \sigma) \leq 1$.
- $F(\varrho, \sigma) = 1$ if and only if $\varrho = \sigma$.
- $F(\varrho, \sigma) = 0$ if ϱ and σ live on orthogonal subspaces.
- Symmetric $F(\varrho, \sigma) = F(\sigma, \varrho)$.
- Let us check consistency. If $\rho = |\Psi\rangle\langle\Psi|$ then $\sqrt{\varrho} = \varrho = |\Psi\rangle\langle\Psi|$. Then,

$$F(\varrho, \sigma) = \text{Tr}(\sqrt{|\Psi\rangle\langle\Psi|\sigma|\Psi\rangle\langle\Psi|})^2 = \langle\Psi|\sigma|\Psi\rangle \text{Tr}(\sqrt{|\Psi\rangle\langle\Psi|})^2 = \langle\Psi|\sigma|\Psi\rangle. \quad (77)$$

Hence, we got back the formula for the simpler case.

- Defining the Fidelity with a maximum over purifications

$$F(\varrho, \sigma) = \max_{|\Psi_\sigma\rangle} |\langle \Psi_\varrho | \Psi_\sigma \rangle|^2. \quad (78)$$